

PRO VAC[®] 4 ACL[™]

Fort Dodge

Bursal Disease-Newcastle Disease-Bronchitis-Reovirus Vaccine

Mass. Type, Killed Virus

For Veterinary Use Only

For the revaccination of healthy chickens as an aid in the prevention of the signs and lesions associated with infectious bursal disease caused by standard and variant strains, Newcastle disease, infectious bronchitis, Massachusetts type and avian reovirus infections which cause malabsorption syndrome. Progeny of vaccinates are aided in the prevention of signs and lesions associated with infectious bursal disease caused by standard and variant strains, Newcastle disease and reovirus disease via maternal antibodies.

DOSE: Inject 0.5 mL (0.5 cc) intramuscularly or subcutaneously (in the lower neck region) using aseptic technique. Vaccinate only healthy birds. For primed birds, a single dose between 16 and 22 weeks (approximately 4 weeks prior to lay) is indicated.

CAUTION: Store in the dark at 36° to 45°F (2° to 7°C). Do not freeze. Warm to 72°F (22°C) and shake well before using. Do not vaccinate within 42 days before slaughter. Use entire contents when first opened.

Gentamicin and amphotericin B added as preservatives.

WARNING: In case of accidental human injection seek immediate medical attention.

Fort Dodge Animal Health

Fort Dodge, Iowa 50501 USA

U.S. Vet. License No. 112

List No.			
17931	500 mL	1,000 Doses	10151B

NAC No.: 10031641